



# Savate™

## MITICIDE/INSECTICIDE

For Greenhouse and Nursery Use

**ACTIVE INGREDIENT:**

Spiromesifen: 2-oxo-3-(2,4,6-trimethylphenyl) -1-oxaspiro[4.4]non-3-en-4-yl 3, 3-dimethylbutanoate.....45.2%

**OTHER INGREDIENTS:**..... 54.8%

**TOTAL:** 100.00%

Savate contains 4 pounds of spiromesifen per US gallon (480 grams ai per liter).

EPA Reg. No. 432-1280

**STOP - Read the label before use**  
**Keep out of reach of children**  
**CAUTION**

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577  
 For PRODUCT USE Information Call 1-800-334-9745

**Specimen Label**

432-1280 160405AV1

<b>FIRST AID</b>	
<b>If inhaled:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<p><b>NOTE TO PHYSICIAN:</b> No specific antidote is known. Treat symptomatically. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.</p>	

### **PRECAUTIONARY STATEMENTS**

#### **HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

##### **CAUTION**

Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

#### **PERSONAL PROTECTIVE EQUIPMENT (PPE)**

##### **Mixers, loaders and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical resistant gloves (such as natural rubber, EPA selection category A)
- Shoes plus socks

##### **Applicators must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### **USER SAFETY RECOMMENDATIONS**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove Personal Protective Equipment immediately after handling this product.
- Wash the outside of gloves before removing.
- As soon as possible, wash thoroughly and change into clean clothing.

#### **ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish and aquatic invertebrates. Avoid contamination of surface water through runoff or spray drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwater.

#### **SURFACE WATER ADVISORY**

This product may contaminate water through runoff or through drift of spray in wind. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

### **DIRECTIONS FOR USE**

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

**IMPORTANT:** Read entire label and Conditions of Sale before using this product.

### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval **(REI) of 12 hours** following application. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, are:

- Coveralls worn over short sleeved shirt and short pants
- Chemical resistant gloves (such as natural rubber, EPA selection category A)
- Shoes plus socks

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**STORAGE:** Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL: Non-refillable container.** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal.

Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Offer for recycling, if available or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. If burned, stay out of smoke.

## GENERAL INFORMATION

Savate™ is a Flowable (Suspension Concentrate) formulation that is intended for control of mites and whiteflies on ornamental plants, flowers and foliage plants in the greenhouse and field and container nurseries. The active ingredient contained in Savate is active on all mite development stages. However, mite juvenile stages are often more susceptible than adults. Application should be timed to coincide with early threshold levels in a developing mite population. Savate is also highly effective against whitefly nymphs, plus it has a significant effect on the otherwise difficult to control pupal stage. For best results against whitefly, applications should target these life stages.

Savate can be applied by ground equipment. Thorough coverage of all plant parts is required for optimum performance. Performance evaluations should be made 4 – 10 days after application.

Apply specific dosage of Savate as needed for control. For best results good coverage of the upper and lower leaf surfaces is recommended. An adjuvant may be used to improve coverage on hard-to-wet foliage. For best results the treatment should be made when whitefly or mite population begins to build and before a damaging population becomes established. Rate range is provided and is generally dependent on size of the plant and density of the foliage. Apply with ground application equipment in adequate water for uniform coverage.

## PRODUCT FEATURES

- Controls all stages of whitefly (nymphs, pupae, and adults).
- Provides knockdown and residual control of all developmental stages of mites

PESTS CONTROLLED	USE RATE
Spider mites (including twospotted spider mite, southern red mite, lewis mite, tumid mite, maple spider mite, spruce spider mite, honeylocust spider mite, euonymus mite, boxwood spider mite) Tarsonemid mites (including broad mite, cyclamen mite) Tenuipalpid mites (including false spider mite) Rust and blister mites (family <i>Eriophyidae</i> )	1 to 4 fl oz (30 – 120 mL)/100 gallons of spray solution or 1.5 – 6 mL/5 gallons of spray solution
Whiteflies (Including greenhouse, silverleaf and sweetpotato whitefly)	2 to 4 fl oz (60-120 mL)/100 gallons of spray solution or 3-6 mL/5 gallons of spray solution
<b>WHERE TO APPLY</b>	<ul style="list-style-type: none"> <li>• To shrubs, trees (including non-bearing fruit and nut trees), flowers and foliage plants in greenhouses, shade houses and plants grown in field and container nurseries.</li> </ul>
<b>HOW TO APPLY</b>	<ul style="list-style-type: none"> <li>• Apply specific dosage of Savate as needed for control. For best results, apply to achieve uniform coverage of the upper and lower leaf surfaces but avoid excess application that results in runoff from plant foliage.</li> <li>• An adjuvant may be used to improve coverage on hard-to-wet foliage. For best results the treatment should be made when the whitefly or mite population begins to build and before a damaging population becomes established</li> <li>• Savate is effective against the egg and nymphal stages of whiteflies and mites. Control should be directed at these stages.</li> <li>• Apply as a full coverage spray to the foliage using a minimum volume of 1 - 2 qts of final solution per 100 sq ft (or 100 to 200 gal per acre). Actual spray volume will vary depending on the size of plants.</li> <li>• For maximum efficacy against adult whiteflies, use the 4 fl oz rate of Savate. For heavy infestations, two applications may be necessary to control adult whiteflies.</li> <li>• Rate range is provided and is generally dependent on size of the plant and density of the foliage. Apply when pests first appear and prior to leaf damage or discoloration. Apply with ground application equipment in adequate water for uniform coverage.</li> <li>• Do not apply via chemigation.</li> <li>• Do not apply Savate more than three times per season.</li> </ul>

(continued)

(continued)

PESTS CONTROLLED	USE RATE
<b>RESISTANCE MANAGEMENT</b>	<ul style="list-style-type: none"><li>• Savate contains an active ingredient with a novel mode of action. Studies to determine cross-resistance with Savate linked to other commercial insecticide/ miticides have demonstrated no cross-resistance. However, repeated use of any plant protection product may increase the development of resistant strains of pests, including insects and mites.</li><li>• Using Savate in successive miticide applications during the same cropping cycle is not recommended. Use Savate as a part of a sound resistance management program that includes rotation with other treatments with different modes of action.</li></ul>
<b>PLANT TOLERANCE</b>	<ul style="list-style-type: none"><li>• Savate has been evaluated for phytotoxicity on a wide range of ornamental plants. However, due to the large number of species and varieties of ornamental plants, it is impossible to test every one for tolerance to Savate. The professional user should determine if Savate could be used safely prior to commercial use. In a small area, test the recommended rates on a small number of plants for phytotoxicity prior to widespread use. Before using Savate in tank mixture with other products and adjuvants, test the mixture on a small number of plants for phytotoxicity prior to widespread use.</li><li>• Do not use Savate on geraniums (<i>Pelargonium</i> sp.), <i>Peperomia</i>, <i>Dracaena</i>, and 'Classy', 'Attache', or 'Vogue' varieties of rose.</li><li>• Insufficient information on tolerance is available for the following ornamentals: lily-of-the-incas (<i>Astromoeria</i>), New Guinea impatiens, <i>Bacopa</i>, ferns, phlox, English ivy (<i>Hedera helix</i>), cyclamen, <i>Agyranthemum</i>, hydrangea, schefflera, <i>Matthiola (stock)</i>, Mexican heather, <i>Lobelia</i>, fuschia, <i>Cordyline</i>, croton, neanthebella palm, and primula. It is recommended that Savate not be used on these plants.</li><li>• Do not apply more than 2 fl oz per 100 gal per application or 4 fl oz per cropping cycle to chrysanthemum, Shasta daisy, snapdragon, impatiens, verbena, lantana, gerbera daisy, or marigolds.</li></ul>
<b>SPRAY DRIFT</b>	<ul style="list-style-type: none"><li>• Avoid drift or runoff into lakes, reservoirs, rivers, permanent streams, marshes, potholes, vernal pools, natural ponds, estuaries, or commercial fish farm ponds. Do not apply when winds are greater than 15 mph and avoid gusty conditions.</li></ul>
<b>IRRIGATION TIMING</b>	<ul style="list-style-type: none"><li>• If irrigation is used, conduct irrigation efficiently to prevent excessive loss of irrigation water through runoff. Time applications to allow sprays to dry prior to rain or sprinkler irrigation. Allow at least 24 hours between applications of product and any irrigation that results in surface water runoff into lakes, reservoirs, rivers, permanent streams, marshes, potholes, vernal pools, natural ponds, estuaries, or commercial fish farm ponds.</li></ul>
<b>ENDANGERED SPECIES</b>	<ul style="list-style-type: none"><li>• The use of any pesticide in a manner that may kill or otherwise harm endangered species or adversely modify their habitat is a violation of Federal law.</li></ul>
<b>RESTRICTIONS</b>	<ul style="list-style-type: none"><li>• Do not use through irrigation systems.</li><li>• Do not use on vegetable gardens.</li><li>• Do not use on plants intended for use as feed or forage.</li><li>• Do not use on bearing fruit or nut trees.</li></ul>
<b>PACKAGE HANDLING</b>	<ul style="list-style-type: none"><li>• Savate is packaged in polyethylene containers. Do not allow product or containers to freeze.</li></ul>

<b>COMPATIBILITY / MIXING / ORDER-OF-MIXING</b>
<ul style="list-style-type: none"><li>• Savate is physically and biologically compatible with many registered pesticides and fertilizers or micronutrients. When considering mixing Savate with other pesticides, or other additives, first contact your supplier for advice. For further information, contact your local Bayer representative. If your supplier and Bayer representative have no experience with the combination you are considering, you should conduct a test to determine physical compatibility. To determine physical compatibility, add the recommended proportions of each chemical with the same proportion of water as will be present in the chemical supply tank, into a suitable container, mix thoroughly and allow to stand for five minutes. If the combination remains mixed, or can be readily remixed, the mixture is considered physically compatible.</li><li>• Savate may be used with other recommended pesticides, fertilizers and micronutrients. The proper mixing procedure for Savate alone or in tank mix combinations with other pesticides is:<ol style="list-style-type: none"><li>1) Fill the spray tank 1/4 to 1/3 full with clean water;</li><li>2) While recirculating and with the agitator running, add any products in PVA bag. Allow time for thorough mixing;</li><li>3) Continue to fill spray tank with water until 1/2 full;</li><li>4) Add any other wettable powder (WP) or wettable granules (WG) products;</li><li>5) Add the required amount of Savate, and any other "flowable" (FL or SC) type products;</li><li>6) Allow enough time for thorough mixing of each product added to tank;</li><li>7) If applicable, add any remaining tank mix components: emulsifiable concentrates (EC), fertilizers and micronutrients.</li><li>8) Fill spray tank to desired level and maintain constant agitation to ensure uniformity of spray mixture.</li></ol></li><li>• Mix pesticides in areas not prone to runoff such as concrete mixing/loading pads, disked soil in flat terrain or graveled mix pads, or use a suitable method to contain spills and/or rinsate. Properly empty and triple-rinse pesticide containers at time of use.</li></ul>

**For Use in California:**

Allow growth of a vegetative filter strip within 25-ft (on which this product should not be applied) along lakes, reservoirs, rivers, permanent streams, marshes, potholes, vernal pools, natural ponds, estuaries or commercial fish farm ponds. This requirement does not apply to retention or holding ponds used to collect and recycle water for irrigation.

### **IMPORTANT: READ BEFORE USE**

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of liability before using this product.

If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following conditions, disclaimer of warranties and limitations of liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, because of manner of use and other factors beyond Bayer CropScience LP's control it is impossible for Bayer CropScience LP to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Bayer CropScience LP disclaims any liability whatsoever for special, incidental or consequential damages, resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.

Bayer (reg'd), the Bayer Cross (reg'd) and Savate™ are trademarks of Bayer.

Produced for:  
Bayer Environmental Science  
A Division of Bayer CropScience LP  
2 T. W. Alexander Drive  
Research Triangle Park, NC 27709

Bayer